

Annual Sedimentation Rates and Role of the
Resuspension Processes Along a Vertical Cliff
(Ligurian Sea, Italy)

Giorgio Bavestrello†, Riccardo Cattaneo-Vietti†, Carlo Cerrano†,
Roberto Danovaro‡ and Mauro Fabiano††

†Istituto di Zoologia
Università di Genova
Via Balbi 5
I-16126 Genova, Italy

‡Istituto di Scienze Ambientali
Marine
Università di Genova
Corso Rainusso 14
S. Margherita L.
I-16038, Genova, Italy

††Dipartimento di Biologia
Animale ed Ecologia
Università di Cagliari
Viale Poetto 1
I-09100 Cagliari, Italy



ABSTRACT

BAVESTRELLO, G.; CATTANEO-VIETTI, R.; CERRANO, C.; DANOVARO, R., and FABIANO, M., 1994. Annual sedimentation rates and role of the resuspension processes along a vertical cliff (Ligurian Sea, Italy). *Journal of Coastal Research*, 11(3), 690-696. Fort Lauderdale (Florida), ISSN 0749-0208.

The annual sedimentation rates along a Portofino Promontory (Ligurian Sea, Mediterranean) cliff were studied using small sediment traps fixed directly to the rocks at 15, 20 and 25 m depth.

Monthly comparisons were made of organic and inorganic matter originating from biological production and cliff erosion in the upper water levels, with suspended matter collected in the water column close to the bottom. The highest quantity of coarse matter, related to the rainfall, was collected in the top trap. Fine sediments were due to the local sea conditions, i.e., wave suspension, and increased from the shallow to the deepest trap. This suggests that resuspension of fine sediments from the sea floor may represent an important fraction of settling matter at lower levels of the submerged coastal cliff. TOM and TSM in the water column are well correlated both at the interface and 1 m above sea floor but not with the amounts of sedimented material. Their annual trends reflect rainfall values, suggesting an influence of the terrigenous outflows.

ADDITIONAL INDEX WORDS: Sediment trap, cliff erosion, coastal sediments, suspended matter.

INTRODUCTION

In recent years, several studies have been carried out on the coupling between pelagic primary production and the structure and functioning of the soft-bottom communities (HARGRAVE *et al.*, 1976; SMETACEK, 1980; TAGUCHI, 1983; BHOSLE *et al.*, 1989). Along coastal cliffs, little is known about this relationship (HEDGES *et al.*, 1988; EVANS *et al.*, 1980; GULLISKEN, 1982; BAVESTRELLO *et al.*, 1991) which appears to be affected not only by the pelagic primary production and local hydrographic regime (SMETACEK, 1980), but also by other factors, such as rainfall and wave action (BAVESTRELLO *et al.*, 1991). Only a small fraction of the primary production is used by hard bottom communities inhabiting the vertical cliff, while a large quantity of organic matter falls on the sea floor, playing an important role in the trophic chains of soft-bottom communities (ODUM and DE

LA CRUZ, 1967; MANN, 1972; HEDGES, *et al.*, 1988; PELET, 1977; EVANS *et al.*, 1980). During rough sea periods, this sedimented matter is resuspended by wave action and contributes to the organic fraction of the water column (ZUNINI-SERTORIO and FABIANO, 1991).

In the present study, the nature and origin of the settling matter and its contribution to resuspended sediments were investigated using traps directly fixed at different depths along a high-energy cliff of Portofino Promontory (Ligurian Sea). In order to understand which factors influence the sedimentation processes, data collected from the traps were compared to the seasonal cycles of the total organic matter (TOM) and total suspended matter (TSM) collected at the water-sediment interface and, in the water column, 1 m above the bottom.

METHODS

Three sediment traps have been placed adherent to a vertical cliff near Punta del Faro (Por-




Figure 1. Sampling area at Portofino Promontory (Ligurian Sea): Punta del Faro (a). Paraggi Bay (b) is also marked.

tofino Promontory, Ligurian Sea, Mediterranean Sea, Figure 1). This kind of trap consists of a rectangular funnel with a mouth of 2 cm × 20 cm converging into a cylindrical collector 20 cm long

(BAVESTRELLO *et al.*, 1991). The first trap was at 15 m depth, inside a pre-coralligenous community, characterized by the green alga *Halimeda tuna*; the second was located on a slightly over-

Table 1. Meteorological data and sediment amount in traps and water column.

Date	Average Wave Height (mm/cm)	Average Rain-fall (mm/day)	Trap 1 15 m (g/m ² /day)			Trap 2 20 m (g/m ² /day)			Trap 3 25 m (g/m ² /day)			TOM (mg/l)		TSM (mg/l)	
			cs	fs	ad	cs	fs	ad	cs	fs	ad	in	ab	in	ab
9.2.90	41.3	1.7	0.90	2.56	0.01	1.00	8.18	3.10	1.12	7.28	0.12	2.56	1.66	9.85	7.55
16.3.90	53.5	0.6	1.06	5.55	0.60	0.83	8.33	1.19	0.78	10.60	0.04	2.70	2.83	8.10	9.03
5.5.90	47.0	6.2	2.42	1.15	0.00	1.67	1.67	0.00	1.66	2.72	0.00	3.52	2.8	9.78	6.83
1.6.90	17.0	0.3	0.23	2.08	0.00	0.00	0.98	0.89	0.00	1.99	0.02	4.4	1.98	12.94	5.14
14.6.90	63.6	0.8	0.00	2.58	1.36	2.54	7.77	6.11	0.00	8.51	1.18	3.19	2.15	11.39	9.77
10.7.90	45.6	0.2	0.00	1.68	0.07	1.60	1.31	2.71	1.33	1.67	0.62	2.73	3.09	9.93	10.84
2.8.90	10.0	0.9	1.17	0.96	0.01	2.12	1.17	1.68	1.17	1.00	0.40	4.05	3.41	13.06	9.74
23.8.90	27.5	2.8	2.80	2.42	1.61	4.29	0.91	4.64	2.06	2.37	0.68	3.85	3.17	16.04	10.93
6.9.90	30.0	5.3	2.03	1.79	0.69	1.75	0.48	0.71	1.26	2.05	0.14	6.1	3.87	21.40	7.90
19.9.90	30.7	2.1	2.33	3.50	0.71	5.24	0.88	1.51	4.91	4.51	0.01	5.18	5.66	18.50	17.69
7.11.90	59.9	12.7	10.14	5.63	1.81	2.30	0.86	0.88	8.74	8.85	0.40	4.45	4.35	14.13	15.00
3.12.90	53.9	4.6	6.20	4.16	0.35	5.83	6.66	1.92	7.76	10.03	1.58	4.07	4.05	13.34	10.38
1.2.91	31.1	6.1	2.84	6.55	0.03	6.94	9.64	2.67	2.68	6.79	0.03	4.78	3.7	17.70	10.88

cs = coarse sediment; fs = fine sediment; ad = algal debris; in = interface; ab = 1 m above




Figure 3. Annual trends of the collected sediments in the three traps. A, B, C: fine matter in traps 1, 2, 3, respectively. D, E, F: coarse matter in traps 1, 2, 3, respectively. The data are reported as daily average of the considered period.

observed, corresponding to heavy rainfall and rough sea periods.

Differences arise when studying separately the two granulometric fractions. The fine sediments collected by the superficial trap (Figure 3A) are abundant mainly in two distinct periods as follow: February–first half March (about $6 \text{ g m}^{-2} \text{ d}^{-1}$) and from October to January (about $6\text{--}7 \text{ g m}^{-2} \text{ d}^{-1}$) and are related to rainfall ($n = 13$; $p < 0.05$). In the second and third trap (Figure 3B and C), the fine sediments collected show a strong correlation with the average wave height ($n = 13$; $p < 0.05$ and $p < 0.001$ respectively) with three peaks in February–first half of March, June and October–January (about $8\text{--}10 \text{ g m}^{-2} \text{ d}^{-1}$).

The coarse sediments are mainly composed of inorganic matter derived from the erosion of the cliff and animal debris (exuvias, serpulids tubes, shells). Most of this fraction was collected in the first trap (about $10 \text{ g m}^{-2} \text{ d}^{-1}$) in October (Figure 3D), and its annual trend is again closely linked to the rainfall ($n = 13$; $p < 0.001$). Also the deepest trap (Figure 3F) shows a trend significantly correlated with rainfall ($n = 13$; $p < 0.001$), while no correlation can be found for trap 2, probably as a result of the particular slope of the cliff in this point. Algal debris has been collected mainly at 20 m depth (Table 1), inside the *Dictyota dichotoma* facies.

In the water column, the trends of TOM (Figure

dynamic conditions. This is well documented by the comparison between the present data and those collected by BAVESTRELLO *et al.* (1991) from Paraggi Bay, a locality very close to Punta del Faro (Figure 1), but much more sheltered from wave action with a seven times higher sedimentation rate. At Punta del Faro, in fact, the cliff is exposed to the main littoral ligurian current which flows from east to west and measures about 30 cm sec⁻¹ (STOCCHINO and TESTONI, 1979). The Paraggi Bay, on the contrary, can be considered a decantation area, because a secondary branch of the main current slows down, flowing along the eastern coasts of the Portofino Promontory (SARA' *et al.*, 1978) forming the so-called "Gulf of Tigullio circuit". These different hydrodynamic conditions are not only at the basis of these quantitative differences, but also can define the qualitative composition of the settling matter. At Paraggi Bay, for instance, the fine sediment fraction, well represented during a large part of the year, is related to the coarse sediments owing to a reduced resuspension incidence. The opposite takes place at Punta del Faro, where the annual trends of the two fractions seem to be independent. Indeed, the fine sediment is more directly linked to wave action than the coarse one and is mainly controlled by the rainfall, which washes away the terrigenous detritus into the sea. Even if the organic fraction of the coarse sediment (animal debris) is quantitatively important, it is lighter than the inorganic one and consequently has a minor influence on the weight relationship. Obviously only the heavier fraction (gravel), originating from the erosion of the cliff, is linked to the rainfall. Along the Portofino Promontory cliff, the fine sedimented matter in agreement with GARDNER's (1977) results was related to the sea conditions, also determining an important resuspension. On average, fine sediments in the deepest trap constituted 66.6% of the total settling matter, and only 52.2 and 55.0% in the traps placed at 15 and 20 m depth. If we assume that the top trap was not significantly influenced by resuspended sediments, in the deepest trap a good fraction of fine sediments (12–15% of total sediment collected) could be related to the resuspension process.

Algal debris are also linked to the sea conditions. This fraction was mostly found in the traps during autumn, in connection with the end of the biological cycle of many macroalgae, but no direct relationship can be envisaged between the cycle of the algal debris and that of TSM and TOM in

the water column. Algal debris before contributing to the particulate suspended matter undergo physical and biological processes, *i.e.*, dispersion, degradation, deposit-feeders ingestion, etc. These processes delay the presence of algal debris in the suspended particulate matter.

Finally, TOM and TSM, as expected, are well correlated both at the interface and 1 m above sea floor, but not with the amount of sedimented material. Their annual trends reflect rainfall values, suggesting a terrigenous origin of this kind of material.

LITERATURE CITED

- BAVESTRELLO, G.; CATTANEO-VIETTI, R.; DANOVARO, R., and FABIANO, M., 1991. Detritus rolling down a vertical cliff on the Ligurian Sea (Italy): The ecological role in hard bottom communities. *Pubbl. St. Zool. Napoli. I.—Marine Ecology*, 12, 281–292.
- BHOSLE, N.B.; SAWANT, S.S.; SANKARAN, P.D., and WAGH, A.D., 1989. Sedimentation of particulate material in stratified and nonstratified water columns in the Bombay High area of the Arabian Sea. *Marine Ecology Progress Series*, 57, 225–236.
- EVANS, R.A.; GULLIKSEN, B., and SANDNES, O.K., 1980. The effect of sedimentation on rocky bottom organisms in Balsfjord, Northern Norway. In: FREELAND, H.J.; FARMER, D.M., and LEVINGS, C.D. (eds.), *Fjord Oceanography*. Plenum, New York, p. 603–607.
- GARDNER, W.D., 1977. Fluxes, Dynamics and Chemistry of Particulates in the Ocean. Ph.D. Thesis. MIT/WHOI Joint Program in Oceanography, 405p.
- GIANMARINO, S.; NOSENGO, S., and VANNUCCI, S., 1959. Risultanze geologico-paleontologiche sul conglomerato di Portofino (Liguria orientale). *Atti Istituto Geologia, Università di Genova*, 7, 305–363.
- GULLIKSEN, B., 1982. Sedimentation close to a near vertical rocky wall in Balsfjorden, Northern Norway. *Sarsia*, 67, 21–27.
- HARGAVE, B.T. and BURNS, M.M., 1976. Assessment of sediment trap collection efficiency. *Limnology and Oceanography*, 24, 1124–1136.
- HARGRAVE, B.T.; PHILLIPS, G.A., and TAGUCHI, S., 1976. Sedimentation measurements in Bedford Basin, 1973–1974. *Technical Report of Marine Service, Environment (Canada)* 608, 1–147.
- HEDGES, J.I.; CLARK, W.A., and COWIE, G.L., 1988. Organic matter sources to the water column and superficial sediments of a marine bay. *Limnology and Oceanography*, 33(5), 1116–1136.
- LUND-HANSEN, L.C., 1991. Sedimentation and sediment accumulation rates in a low-energy embayment. *Journal of Coastal Research*, 7, 969–980.
- MANN, K.H., 1972. Macrophyte production and detritus chains in coastal waters. *Memorie Istituto di Idrobiologia, Pallanza*, 29 suppl., 353–383.
- MARSH, J.B. and WEINSTEIN, W.J., 1966. A simple charring method for determination of lipids. *Journal of Lipid Research*, 7, 574–576.
- ODUM, E.P. and DE LA CRUZ, G., 1967. Particulate organic detritus in a Georgia salt marsh estuarine ecosystem.